


The simplified Pentacel® and Quadracel® immunization series incorporates A VACCINATION SCHEDULE WITH FEWER INJECTIONS¹⁻³

CDC^a schedule through 6 years of age³

Incorporation of Pentacel and Quadracel vaccines on the immunization schedule



^a CDC = Centers for Disease Control and Prevention; ^b Depending on the type of vaccine used, a 3rd dose may be required³;

^c For certain high-risk groups, a dose at 6-11 months of age is recommended.³

INDICATION FOR PENTACEL AND QUADRACEL VACCINES

Pentacel vaccine is indicated for active immunization against diphtheria, tetanus, pertussis, poliomyelitis, and invasive disease due to *H influenzae* type b.

Pentacel vaccine is approved for use as a 4-dose series in children 6 weeks through 4 years of age (prior to fifth birthday).



Quadracel vaccine is indicated for active immunization against diphtheria, tetanus, pertussis, and poliomyelitis. A single dose of Quadracel vaccine is approved for use in children 4 through 6 years of age as a fifth dose in the diphtheria, tetanus, pertussis vaccination (DTaP) series, and as a fourth or fifth dose in the inactivated poliovirus vaccination (IPV) series, in children who have received 4 doses of Pentacel vaccine and/or DAPTACEL® (Diphtheria and Tetanus Toxoids and Acellular Pertussis Vaccine Adsorbed).

IMPORTANT SAFETY INFORMATION FOR PENTACEL AND QUADRACEL VACCINES

Contraindications to vaccination with Pentacel or Quadracel vaccine include: a severe allergic reaction (eg, anaphylaxis) to any ingredient of the vaccine, or following any other diphtheria toxoid-, tetanus toxoid-, pertussis antigen-containing vaccine, inactivated poliovirus vaccine, or *Haemophilus influenzae* type b vaccine (Pentacel vaccine only); encephalopathy within 7 days after a previous dose of a pertussis antigen-containing vaccine with no other identifiable cause; or a progressive neurologic disorder.

Please see additional Important Safety Information on the following page.

Please see full Prescribing Information for [Pentacel](#) or [Quadracel](#) vaccine.



IMPORTANT SAFETY INFORMATION FOR PENTACEL AND QUADRACEL VACCINES (Continued)

Carefully consider benefits and risks before administering Pentacel or Quadracel vaccine to persons with a history of: fever $\geq 105^{\circ}\text{F}$, hypotonic-hyproresponsive episode (HHE), or persistent, inconsolable crying lasting ≥ 3 hours within 48 hours after a previous pertussis antigen-containing vaccine; seizures within 3 days after a previous pertussis antigen-containing vaccine; Guillain-Barré syndrome occurring within 6 weeks following receipt of a prior vaccine containing tetanus toxoid; or adverse events after a previous dose of Pentacel or Quadracel vaccine or receipt of any other tetanus toxoid-, diphtheria toxoid-, or pertussis antigen-containing vaccine.

For infants and children with a history of previous seizures, an antipyretic may be administered (in the dosage recommended in its prescribing information) at the time of vaccination with Pentacel vaccine and for the next 24 hours.

Apnea following intramuscular vaccination has been observed in some infants born prematurely.

The most common local and systemic adverse reactions to Pentacel vaccine include erythema, swelling, and tenderness at the injection site; fever, fussiness, and abnormal crying. The most common local and systemic adverse reactions to Quadracel vaccine include pain, erythema, and edema at the injection site; myalgia, malaise, and headache. Other adverse reactions may occur.

Vaccination with Pentacel or Quadracel vaccine may not protect all individuals.

Before administration, please see full Prescribing Information for Pentacel or Quadracel vaccine.

References: 1. Pentacel vaccine [Prescribing Information]. Swiftwater, PA: Sanofi Pasteur Inc. 2. Quadracel vaccine [Prescribing Information]. Swiftwater, PA: Sanofi Pasteur Inc. 3. Centers for Disease Control and Prevention. Recommended immunization schedule for children and adolescents aged 18 years or younger, United States, 2019. <https://www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-child-combined-schedule.pdf>. Accessed June 27, 2019.

CPT^{®f} Code for Pentacel vaccine: 90698

CPT Code for Quadracel vaccine: 90696

^f CPT (Current Procedural Terminology) is a registered trademark of the American Medical Association.

Pentacel vaccine is manufactured by Sanofi Pasteur Limited and Sanofi Pasteur SA and distributed by Sanofi Pasteur Inc.

Quadracel vaccine is manufactured by Sanofi Pasteur Limited and distributed by Sanofi Pasteur Inc.

